



CLINICAL CHARACTERISTICS OF CHILDREN UNDER INVESTIGATION FOR DENTAL CAVITIES AMONG CHILDREN

Bukhara State Medical Institute named after Abu Ali ibn Sino, Uzbekistan

Samadova Shahzoda Isakovna

PhD, docent

samadova.shahzoda@bsmi.uz

Abstract: In modern dentistry, despite the use of various methods of prevention and treatment of dental hard tissue caries, it remains one of the most common diseases of dental hard tissue. There are hundreds of reasons that contribute to the development of caries in milk teeth. The most important of them are socio-demographic, biological, dietary, hygienic and nutritional factors. In this case, the occurrence or attenuation of a number of conditions will later determine the early development of caries in children.

Key words: caries disease, dental hard tissues, dental care, epidemiological

Introduction

On a world scale, research is being conducted on the development of effective methods of diagnosis, treatment and prevention of caries among children, providing dental care to patients. Dental hard tissue caries is common among the population of different geographical regions and different countries of the world. (1). Due to the prevalence of dental caries in children and the low effectiveness of treatment and prevention measures, it remains a serious health problem. (2)

In the scientific research of a number of local and foreign scientists on the diagnosis, treatment and prevention of caries of dental hard tissues in children, the most frequent referral to specialists with the diagnosis of caries (5) and as the main factors of the occurrence of the pathological process - socio-demographic, biological, hygienic and type of nutrition (6), as well as mental - they showed that the disturbance of the physical condition is also an important etiological factor of the development of caries (7).

It is known that the creation of methods that can clearly show signs of pathology in research, including clinical-functional, biochemical, immuno-microbiological, social and equipment research (4) results in It is noted that there is a continuous relationship between the demineralization of the hard tissue of the tooth, the change in the mineral content of the oral fluid, the change in the microflora of the tooth-gum pocket, and pathology.

It's the last year In our country, in the prevention and treatment of dental caries in children with cerebral palsy, ROS S gel (1) is used, and for the treatment of primary caries in children with rheumatic diseases, Tooth Mousse is made from Nanofluor gel, a new generation representative of bioactive fluorine storage. (7) is recommended to use.

In addition, physical and functional discomfort A program aimed at improving the quality of life (9) was recommended in cases of disorders, changes in the child's emotional, social and family stability, which led to the creation of new methods of prevention and treatment. A disease in the deterioration of the hygienic condition of the oral cavity of children development x amnesty and degree of severity are quite high, and such factors influence the development and clinical course of the disease in many ways (7).



In addition, the dental hard tissues among different strata of the population a number of scientists have conducted scientific research on early diagnosis, assessment of risk factors and increasing the effectiveness of therapeutic treatment (9). However, treatment methods aimed at normalizing the cariogenic microflora along with the remineralization of dental hard tissues in children of different age groups have not been improved and have not been applied to dental practice .

The purpose of the study. prevention of dental caries in children and justification of complex treatment methods.

Research material and method. Based on the results of annual statistical reports and two epidemiological statistical reviews (2018-2023) as the basis of the research, a collective study of the spread of dental caries and its intensity among the population of Bukhara region, an analysis of the direct and indirect results of its implementation, and the development of individual preventive programs of dental caries are chosen. received. In the implementation of the set goal:

Monitoring of prevalence and intensity of dental caries in the population of cities and districts of Bukhara region during a 3-year period; - development of an individual preventive program and evaluation of its clinical effectiveness after its introduction among children and adolescents was carried out.

In order to determine the trends in the dynamics and speed of dental caries prevalence in the city and district, the average indicators of these indicators in the 3 main (6-9, 10-12, 13-15) age categories of the population living in cities and rural areas were calculated and the reliability of the differences over a 3-year period was evaluated.

Examinations were carried out at the Department of Pediatric Dentistry of the Bukhara State Medical Institute together with specialists trained in assessment methods according to the criteria of the regional WHO (WHO, 1997).

The results of the examination were entered into single record cards and analyzed with the help of a special computer program, which made it possible to create a database of statistical analysis. A total of 4181 people in 2018 and 4955 people in 2020 were analyzed. Prevention of dental caries in children of different age groups and the fulfillment of the assigned tasks in the period from 2017 to 2019, the dental clinic of the Bukhara State Medical Institute (head doctor - M.A. Astanov) and the department of pediatric dentistry (the head of the department, Ph.D., Assoc. Kamalova) for 3 years, a comprehensive examination of 134 clients on their dental status, assessment of the need for treatment of children's caries, and an individualized program based on rational approaches to its implementation in preschool children were developed. Comparative evaluation of KPO indicators was carried out through an additional control group, which included 36 children aged 6 to 15 years who had not previously participated in preventive programs.

Results of research work. Main dental diseases in children among the population of Bukhara region in 2019 - the average indicators of prevalence and intensity of dental caries and periodontal inflammatory diseases are shown in caries among early and middle-aged children of Bukhara region were formed, including 84% of children under six years of age have dental caries, and 75% of 12-year-old children have dental caries. Almost half of the 15-year-olds had periodontal inflammation..

Children between The analysis of disease dynamics showed the stability of the average indicators of the intensity of caries of temporary teeth in 6-year-old children during the 3 - year



period. There was a redistribution of components in the KPO: the number of carious teeth decreased by 15%, the number of filled teeth increased by 2.5 times compared to 2018. In 2020, it was noted that the gradation of dental caries development in 12-year-old children is low (WHO). The development of caries of dental hard tissue in patients belonging to each age category decreased by 30%. The total number of tooth extraction procedures in 2020 has been reported to decrease by 50.65%. This, of course, indicates the development of clinical diagnosis and treatment methodology in dental practice.

A certain decrease in the prevalence of signs of damage to periodontal tissues in early-age children can be seen in this diagram. In particular, in the last two years (compared to 2020-2021 and 2018-2019), the prevalence of periodontal tissue damage has decreased by 15 and 16.5 percent, respectively.

The rate of dental caries in 12-year-old children living in the city was higher than in children living in rural areas, but the value of the "K" component was found to be almost the same in urban and rural children. The number of filled teeth ($p < 0.001$) was more in children living in the city (1.32 ± 0.07) than in children living in the countryside (0.39 ± 0.04). Among 15-year-olds, the value of KPO indices was higher in urban than in rural districts (4.30 ± 0.21 and 2.48 ± 0.15 , $p < 0.001$). Also, the values of "K" and "P" components (1.84 ± 0.10 and 2.26 ± 0.11) are higher in urban than in rural children (1.43 ± 0.07 and 0.72 ± 0.06). ($p < 0.001$) was.

Thus, in the analysis of dental caries rate, rural children showed a trend towards higher rates of caries in both temporary teeth and permanent teeth than their urban counterparts.

Summary. For the purpose of complex dental treatment, the filling of cavities of milk teeth (fissure sealing), the appointment of vitamin D 3 and calcium preparations (under the supervision of a pediatrician), the exogenous application of fluoride varnish, training in personal hygiene of the oral cavity, and repeated courses of the treatment every 6-8 months are necessary. became the basis for preventing caries of the tissue. The values of the KPO index in 12-year-old children did not differ significantly, although the value of the K component was 2 times higher in rural children. In 15-year old children, the average value of KPU index (3.31 ± 0.15) and its component "K" (1.85 ± 0.10) turned out to be higher ($p < 0.05$). compared to children from rural areas. Urban children (1.69 ± 0.08) were higher than their urban peers (2.80 ± 0.12 and 1.06 ± 0.06 , respectively), and the P component was higher in rural children (1.38 ± 0.07), ($p < 0.05$).

Literature

1. Samadova Shahzoda Isakovna CLINICAL CHARACTERISTICS OF CHILDREN UNDER INVESTIGATION FOR DENTAL CAVIES AMONG CHILDREN. International Journal of Health Systems and Medical Science ISSN: 2833-7433 № 6 June – 2023 P.64-68
2. Samadova Shahzoda Isakovna THE LEVEL OF OCCURRENCE OF DISEASES OF THE MUCOUS MEMBRANE OF THE ORAL CAVITY AND DENTAL CAVIES (literature review) Vol. 22 No. 8 (2023): ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ Выпуск журнала № 22 Часть-8 Р.101-106
3. Samadova Shahzoda Isakovna MODERN METHODS OF DIAGNOSTICS OF LICHEN PLANE European Journal of Modern Medicine and Practice (EJMMP) Vol. 3 No. 4 (2023): EJMMP PUBLISHED: 2023-04-04 P.33-36
4. Самадова Шахзода Исаковна СОВРЕМЕННЫЕ МЕТОДЫ ДИАГНОСТИКИ КРАСНОГО ПЛОСКОГО ЛИШАЯ Journal of new century innovations | www.newjournal.org | Volume27 | Issue-3 Vol. 27 No. 3 (2023) P.118-123



Western European Journal of Modern Experiments and Scientific Methods

Volume 3, Issue 6, June 2025

<https://westerneuropeanstudies.com/index.php/1>

ISSN (E): 2942-1896

Open Access | Peer Reviewed



This article/work is licensed under CC Attribution-Non-Commercial 4.0

5. Sh.I. Samadova Changes In Biochemical Indicators During the Development of Alveolitis. Journal of Eurasian Medical Research Periodical Volume 35| August 2024 ISSN: 2795-7624 P. 56-59

6. Khojaeva*, Otabek Rajabov, Lola Muhsinova, Dilafruz Hamroyeva, Shahzoda Samadova, and Nozima Kazakova. Increasing the effectiveness of complex influence on the condition of parodontal tissue in patients with diseases of the digestive system and having fixed dentures BIO Web of Conferences 121, 03005 (2024) <https://doi.org/10.1051/bioconf/>

7. Samadova Shaxzoda Isokovna New Methods for Treating Inflamed Dental Alveoli and Characteristics of Osteoplastic Materials Used for Treatment. International Journal of Integrative and Modern Medicine Volume 2, Issue 6, 2024 ISSN: 2995-5319 P.806-809

8. Samadova Sh.I PASTKI JAG MOLYAR TISHLARINI OLINGANDAN KEYIN YUZAGA KELADIGAN ALVEOLITLARNI OLDINI OLISH VA KOMPLEKS DAVOLASH. International scientific journal «MODERN SCIENCE AND RESEARCH» VOLUME 3 ISSUE 6 ISSN: 2181-3906 2024 P.298-302