



ALTERATION IN THE CONVULSIVE EFFECT OF KYNURENINE, CORAZOL AND CAFFEINE UNDER THE ACTION OF BICUCULLINE

Mirzaolimov Mirzahid Mirzavalievich¹

PhD, Namangan State University

E-mail: mirzohid_0421@mail.ru Tel: +998974403343

Niyozov Qahramon Adashaliyevich²

senior lecturer, Department of Physiology, Namangan State University.

E-mail: nambiolog@umail.uz Tel: +998902602197

Abstract: In male mice of two strains SHR and C57BL/6, pretreatment with bicuculline (1 mg/kg, s.c.), an antagonist of GABA-A receptors, potentially the convulsant effect of pentylenetetrazole (40mg/kg, ip.) and caffeine (200mg/kg, ip.) and had no effect on that of L-kynurenine (11 meg, i.c.v.). Data suggest that the convulsant effect of L-kynurenine, an endogenous convulsant from neuroactive tryptophan metabolites, is not related to GABA-A receptors. Other data reported elsewhere (e.g. the antagonism of phenibut and baclofen, agonists of GABA-B receptors, to L-kynurenine) suggest that kynurenine-induced seizures are related to GABA-B receptors.

Keywords: convulsive effect, bicuculline, corazol, kinurenin, GABA-A-B receptors, caffeine.

ИЗМЕНЕНИЕ СУДОРОЖНОГО ЭФФЕКТА КИНУРЕНИНА, КОРАЗОЛА И КОФЕИНА ПОД ДЕЙСТВИЕМ БИКУКУЛЛИНА

Мирзаолимов Мирзохид Мирзавалиевич¹

PhD, Наманганский государственный университет

E-mail: mirzohid_0421@mail.ru Tel: +998974403343

Ниёзов Каҳрамон Адашалиевич²

Наманганский государственный университет

E-mail: nambiolog@umail.uz Tel: +998902602197

Аннотация: У самцов мышей двух штаммов SHR и C57BL / 6, предварительной обработки бикукуллином (1 мг / кг, подкожно), антагонистом рецепторов ГАМК-А, потенциально судорожный эффект пентиленететразола (40 мг / кг, внутрибрюшинно) и кофеина (200 мг / кг, внутрибрюшинно) и не оказывал влияния на L-кинуренин (11 мг, icv). Данные свидетельствуют о том, что судорожный эффект L-кинуренина, эндогенного судороги из нейроактивных метаболитов триптофана, не связан с рецепторами ГАМК-А. Другие данные, представленные в других источниках (например, антагонизм фенибута и баклофена,agonистов рецепторов ГАМК-В к L-кинуренину), свидетельствуют о том, что вызванные кинуренином судороги связаны с рецепторами ГАМК-В.



Ключевые слова: судорожный эффект, бикукуллин, коразол, кинуренин, рецепторы ГАМК-А-В, кофеин

BIKUULLIN TA'SIRIDA KINURENIN, KORAZOL VA KOFEINNING KONVULSIV TA'SIRINING O'ZGARISHI

Mirzaolimov Mirzohid Mirzavaliyevich¹

PhD, Namangan davlat universiteti

E-mail: mirzohid_0421@mail.ru Tel:+998974403343

Niyozov Qahramon Adashaliyevich²

Namangan State University,

Annotatsiya: SHR va C57BL / 6 ikkita shtammining erkak sichqonlarida, bikuullin (1 mg / kg, teri ostiga) yuborish, GABA-A retseptorlari antagonisti, pentilenetetrazolning potensial konvulsiv ta'siri (40 mg / kg, ichkariga) va kofein (200 mg / kg), qorin bo'shlig'ida) va L-kinurenining (11 mg, iv) ta'sir ko'rsatmadi. Dalillar shuni ko'rsatadiki, L-kinureninning konvulsiv ta'siri, neyroaktiv triptofan metabolitlaridan endogen ushlanishi GABA-A retseptorlari bilan bog'liq emas. Boshqa manbalarda keltirilgan boshqa ma'lumotlar (masalan, fenibut va baklofennning antigenizmi, GABA-A retseptorlarining L-kinurenining bo'lgan agonistlari) kinurenin bilan tutilish GABA-B retseptorlari bilan bog'liqligini ko'rsatadi.

Kalit so'zlar: konvulsiv ta'sir, bikukullin, korazol, kinurenin, GABA-AB retseptorlari, kofein.

Introduction

Gamma-aminobutyric acid (GABA) is a major neurotransmitter widely distributed throughout the central nervous system (CNS). Because too much excitation can lead to irritability, restlessness, insomnia, seizures, and movement disorders, it must be balanced with inhibition. GABA – the most important inhibitory neurotransmitter in the brain – provides this inhibition, acting like a “brake” during times of runaway stress. Medications for anxiety, such as benzodiazepines, stimulate GABA receptors and induce relaxation. Either low GABA levels or decreased GABA function in the brain is associated with several psychiatric and neurological disorders, including anxiety, depression, insomnia, and epilepsy. Studies indicate GABA can improve relaxation and enhance sleep [1].

Numerous experimental studies have found that the γ -aminobutyric acid (GABA) system of the brain is important in protecting against arousal and seizures caused by excitatory neurokynurene - the endogenous metabolites of the essential amino acid tryptophan [2]. Of the excitatory neurokynurenes, the most active are L-kynurene (kynurene) and quinolinic acid. Among the receptors involved in the excitatory effects of neurokynurenes, the most studied GABA receptors and excitatory amino acid receptors [2, 3]. When comparing kynurene convulsions with convulsions caused by standard convicts (corazol, caffeine, etc.), it was found that the GABA-B receptors and dopamine (DA) receptor blockade play a leading role in the mechanism of kynurene convulsions [2]. GABA-A receptors seem to be much less important, since even such a universal and powerful anticonvulsant, as diazepam, a typical

agonist of GABA-A receptors, is much less effective in protecting against kynurenine convulsions than GABA-B receptor agonists. Experimental therapy of kynurenine convulsions [4] found that protection from them requires first of all the activation of GABA-ergic processes, for example, with the help of GABA-mimetics, and also the strengthening of DA-ergic processes by isolating DA or its predecessors.

The purpose of the study was to obtain additional information about the role of GABA-A receptors in protection against kynurenine seizures. For the analysis, the selective antagonist of the GABA-A receptor bicuculline was used, which has been widely used in recent years for this purpose [3].

Materials and research methods. The experiments were performed on male mice of two lines: outbred albinos (originating from the SHR line) and black C57BL/6 weighing 20-21 g, obtained from the Rappolovo farm near St. Petersburg. Bicuculline in the form of a solution freshly prepared on distilled water was injected subcutaneously 15 minutes before the convulsant. Solutions of corazol and caffeine were injected intraperitoneally, kynurenine into the cerebral ventricles using a semi-automatic device [2].

All preparations were obtained from "Sigma" (USA). The number of animals with typical clonic-tonic convulsions (within 2 hours after the introduction of convulsants) and the number of dead animals were recorded after 2 and 24 hours. Work performed in the **Laboratory of Psychopharmacology, Bekhterev Psychoneurological Research Institute, St. Petersburg**.

Results and discussion. The results of the experiments showed (table) that the preliminary administration of bicuculline significantly enhances the convulsive effect of corasol and caffeine, but not kynurenine. It should also be emphasized that convulsants are used in equally effective - minimal convulsive doses.

Table

The effect of bicuculline on the convulsive effect of kynurenine, corazol and caffeine

	Preparations			Amount of mice					
	dose			outbred (SHR)			C57BL/6		
	s/c	i/p	i/bv	all with dead by convulsion			all with dead by convulsion		
	mg/kg								
S+ kynurenine	-		12	20	2	0	26	6	2
BCC- kynurenine	1		12	20	3	0	26	12	6
S-corazol	*	40		28	6	0	28	4	0
BCC+ corazol	1	40		28	20*	6*	28	24*	0
S+ caffeine	-	200		30	9	1	24	6	2
BCC+caffeine	1	200		30	19*	1	24	16*	6

Abbreviations: S – saline (control), BCC - bicuculline * $p < 0,05$ (with control). s/c – subcutaneously, i/p – intraperitoneally, – inside the ventricle of the brain

Conclusions. The obtained data give reason to support the proposal [2] that GABA-A receptors, which are crucial in the mechanism of corazol and caffeine convulsions, are not involved in the mechanism of kynurenine convulsions, and mainly associated with GABA-B receptors.

REFERENCES

1. Axmerov R. N. et al. ON THE POSSIBILITY OF UNCOUPLED MITOCHONDRIA IN BROWN FAT OF NEWBORN GUINEA PIGS //Scientific Bulletin of Namangan State University. – 2019. – Т. 1. – №. 9. – С. 49-55.
2. Mirzaolimov M. M., Abdullaev G. R., Abdullayev S. S. ROLE OF A CALORIE-RESTRICTED DIET IN PROLONGING THE LIFESPAN OF AN ORGANISM AND ITS MITOCHONDRIAL MECHANISMS //Scientific Bulletin of Namangan State University. – 2019. – Т. 1. – №. 10. – С. 106-112.
3. Niyazmetov B. et al. UNCOU'LED RESPIRATION IN BIRD MITOCHONDRIA: CONNECTION WITH THERMOGENESIS //Scientific Bulletin of Namangan State University. – 2019. – Т. 1. – №. 2. – С. 100-104.
4. Soliev N., Mirzaolimov M. ACTION OF CALCIUM ON THE CONTENT OF PHOSPHOTYDYLCHOLIN, PHOSPHATYL ETHANOLAMINE AND THEIR LYSOFORMS IN THE RAT LIVER MITOCHONDRIA //Scientific Bulletin of Namangan State University. – 2019. – Т. 1. – №. 3. – С. 69-71.
5. Mirzaolimov M. M. et al. THE METHOD OF SEPARATION OF MITOCHONDRIAS AND DETERMINATION OF PHYSIOLOGICAL AND BIOCHEMICAL CHANGES IN ORGANISMS IN ONTOGENESIS //Scientific Bulletin of Namangan State University. – 2020. – Т. 2. – №. 3. – С. 175-178.
6. Мирзаолимов М. М., Рахимжонович М. А. Г. ВЛИЯНИЕ ПЕРЕКИСНОГО ОКИСЛЕНИЯ ЛИПИДА НА МИТОХОНДРИЮ ПЕЧЕНИ КРЫС ПРИ ПОСТНАТАЛЬНОМ ОНТОГЕНЕЗЕ //INTERNATIONAL JOURNAL OF DISCOURSE ON INNOVATION, INTEGRATION AND EDUCATION. – 2020. – Т. 1. – №. 5. – С. 78-86.
7. Niyazmetov B., Akhmedov R., Mirzaolimov M. UNCOU'LED RESPIRATION IN BIRD MITOCHONDRIA: CONNECTION WITH THERMOGENESIS //Bulletin of Namangan State University: Vol. – 2019. – Т. 1. – №. 2. – С. 18.
8. Бохонова Н. С., Мирзаолимов М. М. ВЛИЯНИЕ КАЛОРИЙНО-ОГРАНИЧЕННОЙ ДИЕТЫ НА ПРОДОЛЖИТЕЛЬНОСТЬ ЖИЗНИ ЖИВОТНЫХ //TA'LIM VA RIVOJLANISH TAHLILI ONLAYN ILMUY JURNALI. – 2022. – С. 139-142.
9. Анваров Ф. Р., Мирзаолимов М. М. ГЕРИАТРИЯ (ОПРЕДЕЛЕНИЕ ПОНЯТИЙ; ЗАДАЧИ, СТОЯЩИЕ ПЕРЕД ЭТИМ НАУКАМ; РАЗДЕЛЫ И ДОСТИЖЕНИЯ) //TA'LIM VA RIVOJLANISH TAHLILI ONLAYN ILMUY JURNALI. – 2022. – С. 326-332.
10. Мирзаолимов М. М. и др. ГЕРИАТРИЯ (ОПРЕДЕЛЕНИЕ ПОНЯТИЙ; ЗАДАЧИ, СТОЯЩИЕ ПЕРЕД ЭТИМ НАУКАМ; РАЗДЕЛЫ И ДОСТИЖЕНИЯ) //TA'LIM VA RIVOJLANISH TAHLILI ONLAYN ILMUY JURNALI. – 2022. – С. 143-149.
11. Таджибаева Г. И., Мирзаолимов М. М. КАЛАМУШЛАРДА СУРУНКАЛИ ЭМОЦИОНАЛ СТРЕСС МОДЕЛИНИ ЯРАТИШ ВА ОРГАНИЗМДАГИ БИОКИМЁВИЙ ЎЗГАРИШЛАРНИ АНИҚЛАШ //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMUY JURNALI. – 2022. – С. 166-170.
12. Mirzavalievich M. M., Adashaliyevich N. Q. Soy Protein, Isoflavones, and Cardiovascular Health //International Journal of Scientific Trends. – 2023. – Т. 2. – №. 4. – С. 10-18.



13. Атаканова С. Д. СТРЕСС ТАЪСИРИДА ОРГАНИЗМ АЪЗО ВА ТИЗИМЛАРИДАГИ ФИЗИОЛОГИК-БИОКИМЁВИЙ ЎЗГАРИШЛАР //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMUY JURNALI. – 2022. – Т. 2. – №. 4. – С. 215-219.
14. Сопиев Ш. К., Рўзибоева С. И., Мирзаолимов М. М. ЕДИНСТВО И ВЗАИМОСВЯЗЬ СТРУКТУРЫ СОРЕВНОВАТЕЛЬНОЙ ДЕЯТЕЛЬНОСТИ И СТРУКТУРЫ ПОДГОТОВЛЕННОСТИ //PEDAGOGS jurnali. – 2022. – Т. 10. – №. 4. – С. 220-228.
15. Таджибаева Г. И., Мирзаолимов М. М. МИТОХОНДРИЯНИНГ ТАРКИБИЙ ТУЗИЛИШИ ВА БАЖАРАДИГАН ВАЗИФАЛАРИ //PEDAGOGS jurnali. – 2022. – Т. 8. – №. 1.
16. Сопиев Ш. К., Рўзибоева С. И., Мирзаолимов М. М. ПОДХОДЫ К РАЗВИТИЮ ДЕТСКО-ЮНОШЕСКОГО СПОРТА И ПОСТРОЕНИЮ МНОГОЛЕТНЕЙ ПОДГОТОВКИ И ПОСТРОЕНИЮ МНОГОЛЕТНЕЙ ПОДГОТОВКИ //PEDAGOGS jurnali. – 2022. – Т. 5. – №. 1. – С. 162-170.
17. Мирзаолимов М. М. и др. ГЕРИАТРИЯ (ОПРЕДЕЛЕНИЕ ПОНЯТИЙ; ЗАДАЧИ, СТОЯЩИЕ ПЕРЕД ЭТИМ НАУКАМ; РАЗДЕЛЫ И ДОСТИЖЕНИЯ) //TA'LIM VA RIVOJLANISH TAHLILI ONLAYN ILMUY JURNALI. – 2022. – С. 143-149.
18. Таджибаева Г. И., Мирзаолимов М. М. КАЛАМУШЛАРДА СУРУНКАЛИ ЭМОЦИОНАЛ СТРЕСС МОДЕЛИНИ ЯРАТИШ ВА ОРГАНИЗМДАГИ БИОКИМЁВИЙ ЎЗГАРИШЛАРНИ АНИҚЛАШ //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMUY JURNALI. – 2022. – С. 166-170.
19. Атаканова С. Д. СТРЕСС ТАЪСИРИДА ОРГАНИЗМ АЪЗО ВА ТИЗИМЛАРИДАГИ ФИЗИОЛОГИК-БИОКИМЁВИЙ ЎЗГАРИШЛАР //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMUY JURNALI. – 2022. – Т. 2. – №. 4. – С. 215-219.
20. Мирзаолимов М. М. и др. КИСЛОРОД ВА АЗОТ ФАОЛ ШАКЛЛАРИ СИНТЕЗИ //PEDAGOGS jurnali. – 2022. – Т. 7. – №. 1. – С. 394-400.
21. Рахматуллаев А. М., Мирзаолимов М. М. РОЛЬ НА-К-НАСОСА. АКТИВНЫЙ ТРАНСПОРТ ИОНОВ КАЛЬЦИЯ И ВОДОРОДА В КЛЕТКЕ //Journal of new century innovations. – 2023. – Т. 43. – №. 1. – С. 8-13.