

# STUDY OF THE EFFECT OF SOME ALKALOIDS ON THE CONTRACTILE ACTIVITY OF AORTIC SMOOTH MUSCLE CELLS

**Mirzayeva Yulduzkhon Takhirjonovna**


Alfraganus University, Tashkent, Uzbekistan,

Institute of Biophysics and Biochemistry, National University of Uzbekistan, Tashkent,  
Uzbekistan

[Ymirzayeva@mail.ru](mailto:Ymirzayeva@mail.ru)

**Key words:** smooth muscle, sarcoplasmic reticulum, aorta, phenylephrine,  $\text{Ca}^{2+}$  channel, 1-O-acetylkarakoline, caffeine.

**Relevance.**  $\text{Ca}^{2+}$  ions play a leading role in regulation of contractile and functional activity of cardiac muscles and smooth muscles of blood vessels, which ensure normal activity of cardiovascular system as a whole [1]. In this regard, disturbances in the regulation of  $\text{Ca}^{2+}$  homeostasis are the main cause of pathogenesis of a number of diseases of the cardiovascular system, including heart failure, arrhythmias, myocardial infarction, hypertension and strokes [2]. Therefore, one of the most urgent tasks of modern biophysics, pharmacology and medicine is to study and characterize the mechanisms of pharmacological regulation of  $\text{Ca}^{2+}$ -homeostasis and  $\text{Ca}^{2+}$ -transporting systems of cardiac and smooth muscles providing its maintenance. The aim of the work was to study the effect of 1-O-acetylkarakoline, a derivative of the diterpenoid alkaloid karakolin, isolated from the plant *Aconitum karakolicum*, on the contractile activity of smooth muscle cells (SMC) of the rat aorta. Karakolin has a pronounced antiarrhythmic effect, which is determined by the presence of specific functionally important groups in its structure [3]. Thus, the replacement of the OH-group at the carbon atom C-1 of the lycoketonin skeleton of Karakolin with an acetyl group (Fig. 1) leads to a significant increase in antiarrhythmic activity of 1-O-acetylkarakoline [4].




**Figure 1. Chemical structure of diterpenoid alkaloids karakoline (A) and acetylkarakoline (B)**

**Research methods and materials.** Isolated preparations of smooth muscle segments of aorta of white mongrel rats (200-250 g) were used in the experiments. The rats were slaughtered by cervical dislocation, the thorax was opened, the aorta was extracted and placed in Krebs physiologic solution of the following composition (mM): NaCl-120.4; KCl-5; NaHCO<sub>3</sub>-15.5; NaH<sub>2</sub>PO<sub>4</sub>-1.2; MgCl<sub>2</sub>-1.2; CaCl<sub>2</sub>-2.5; C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>-11.5; pH 7.4. The aorta was cleared of adipose and connective tissue and cut into segments in the form of rings 3-4 mm wide. The dissected aortic segment was placed in a special experimental chamber of 5 ml volume, where it was fixed between the stem of the electromechanical transducer (FT.03, Grass, USA) and the chamber base using silver wire. The experimental chamber was perfused with oxygenated carbogen (95% O<sub>2</sub>, 5% CO<sub>2</sub>) Krebs solution at a constant temperature of 37°C. Before the experiment, aortic segments were pre-stretched with a 1 g load and washed with saline for 60 minutes to achieve equilibrium. Contractions of aortic preparations were induced by applying PE (1 μM) or hyperkalemik solution (KCl, 50 mM) to the experimental chamber. Calcium-free solutions were also used in the experiments, for which Ca<sup>2+</sup> ions were excluded from the Krebs solution and EGTA (1 mM) was added to bind their traces. The contractile activity of aortic preparations was recorded using a PIIT amplifier (Grass, USA) and an Endim 621.02 recorder (Germany). Statistical data processing was performed using OriginPro 7.5 program (OriginLab Corporation; USA). The contraction amplitude was expressed as % of the maximum contraction (taken as 100%) induced by phenylephrine (PE) or hyperkalemic solution and calculated as the arithmetic mean for 4-5 different experiments. Statistical significance of differences between control and experimental values was determined for a series of data using paired t-test. Values of  $p < 0.05$  indicate statistically significant differences.

**Results and their discussion.** Previously, we found that the diterpenoid alkaloids 1-O-acetylkarakoline and karakoline effectively relax rat aortic preparations precontracted with phenylephrine and hyperkalemic solutions [5]. Therefore, the aim of the present work was to further characterize the mechanism of relaxant action of 1-O-acetylkarakoline.

To further characterize the mechanism of relaxant action of 1-O-acetylkarakoline (1-O-AK), we studied its effects on contractions of rat aortic preparations induced by the selective α-adrenoreceptor agonist phenylephrine, in the development of which Ca<sup>2+</sup> ions entering the SMCs not only through potential-dependent, but also through receptor-regulated Ca<sup>2+</sup> channels, and also released from the sarcoplasmic reticulum (SR) participate [6,7]. In these studies, we found that 1-O-AK and under conditions of PE-induced contracture effectively relaxes rat aortic preparations. The relaxant effect of 1-O-acetylkarakolin was also dose-dependent, and at its concentration of 20 μM it caused relaxation of the aortic preparation by 24.5 ± 3.2%, and maximum relaxation up to 81.1 ± 4.1% was observed at its concentration of 150 μM. The magnitude (concentration 50% of maximal effect) of EC<sub>50</sub> of 1-O-acetylkarakolin under these conditions was 41.9 ± 4% μM. At the same time, the relaxant effect of 1-O-acetylkarakolin, under conditions of PE-induced contracture, was also observed in calcium-free solutions, the development of which is mainly provided by Ca<sup>2+</sup> ions released from SR.


Under these conditions, PE induced contraction of the rat aortic preparation, which was  $70.4 \pm 4.1\%$  less than the contraction induced by PE in the presence of  $\text{Ca}^{2+}$  ions. Addition of 1-O-acetylkarakolin under these conditions resulted in an additional  $23.2 \pm 3.8\%$  reduction in contraction amplitude (Figure 2.). The results of these experiments indicate that the relaxant effect of 1-O-acetylkarakolin, in conditions of PE-induced contracture, is mainly due to its effect on the entry of  $\text{Ca}^{2+}$  ions into SMCs through receptor-controlled  $\text{Ca}^{2+}$  channels of plasmalemma and on the release from SR. It should be noted that the relaxant effect of karakolin in conditions of PE-induced contracture, as well as in conditions of KCL-induced contracture, was less pronounced in comparison with the effect of 1-O-acetylkarakolin.



**Figure 2. Effect of 1-O-acetylkarakoline on contractions of rat aortic preparations induced by phenylephrine.** On the ordinate axis is the strength of aortic contraction expressed as a percentage of the control obtained by the action of 1 μM phenylephrine and taken as 100%. ( $P < 0,05$ ;  $n = 5$ ).

The results of these experiments provide convincing evidence that the relaxant effect of the alkaloid 1-O-AK is realized mainly as a result of its influence on  $\text{Ca}^{2+}$ -transporting systems of SMCs controlled by  $\alpha$ -adrenoreceptors and related intracellular signaling cascades. It is known that stimulation of  $\alpha$ -adrenoreceptors, inositol-1,4,5-triphosphate ( $\text{IP}_3\text{R}$ ), is accompanied by activation of  $\text{Ca}^{2+}$  channels of plasmalemma and sarcoplasmic reticulum [6]. At the same time, the release of  $\text{Ca}^{2+}$  ions from SR via  $\text{IP}_3\text{R}$  contributes mainly to the increase of  $[\text{Ca}^{2+}]_i$  and activation of SMC contraction. Taking this into account, we studied the effect of the investigated alkaloid on the effects of caffeine, which affects the contractile activity of SMCs by specifically modifying  $[\text{Ca}^{2+}]_i$  [8]. In particular, caffeine activating RyR can increase the release of  $\text{Ca}^{2+}$  ions from SR and stimulate the contraction of SMCs [9]. At the same time, caffeine can interact with  $\text{IP}_3\text{R}$  and inhibiting it, suppresses the release of  $\text{Ca}^{2+}$  ions from SR and causes smooth muscle relaxation [10].

Taking into account these peculiarities of caffeine action, we studied the effect of 1-O-AK on contractions of aortic preparations induced by caffeine in the absence of  $\text{Ca}^{2+}$  ions in the incubation medium. As can be seen in Fig. 2, addition of caffeine to the solutions containing  $\text{Ca}^{2+}$  ions induced a contraction of aortic preparations, which was  $67.5 \pm 4.1\%$  of the contraction induced by phenylephrine. When  $\text{Ca}^{2+}$  ions are excluded from the incubation medium, caffeine-induced contraction of aortic preparations decreases to  $33.3 \pm 3.4\%$  of the control obtained in the presence of  $\text{Ca}^{2+}$  ions. Addition of the alkaloid 1-O-AK (150  $\mu\text{M}$ ) under these conditions observed an additional reduction in caffeine-induced contraction to  $21.1 \pm 4.4\%$ , from control (Fig.3).



**Figure 3. Effect of 1-O-acetylkarakoline on contractions of rat aortic preparations induced by caffeine.** On the ordinate axis is the strength of aortic contraction expressed as a percentage of the control obtained by the action of 1  $\mu\text{M}$  phenylephrine and taken as 100%, ( $P < 0,05$ ;  $n=4$ ).

The results of these experiments indicate that 1-O-AK suppresses caffeine-induced contractions of aortic preparations. Taking into account that this effect of the studied alkaloid was detected under the conditions of absence of  $\text{Ca}^{2+}$  ions in the incubation medium, in which their entry into SMCs from the extracellular medium is excluded, we can assume that it is due to the effect of the alkaloid on the release of  $\text{Ca}^{2+}$  ions from SR via RyR or  $\text{IP}_3\text{R}$ .

Thus, analysis of the data obtained in these experiments shows that the alkaloid 1-O-AK effectively relaxes rat aortic preparations precontracted by the  $\alpha$ -adrenoreceptor agonist phenlephrine. Taking into account that contractions induced by this antagonist are mainly caused by activation of  $\text{Ca}^{2+}$  ions release from SR via  $\text{IP}_3\text{R}$ , we can conclude that the relaxant effect of the studied alkaloid is caused by suppression of  $\text{Ca}^{2+}$  ions transport at SR level.

## Literature



1. Fleming I. and R. Busse. "Signal transduction of eNOS activation," *Cardiovascular Research*, vol. 43, no. 3, pp. 532-541, 1999.
2. Jahangirov F.N., Sultankhodjaev M.N., Tashkhodjaev B., Salimov B.T. Diterpenoid alkaloids as a new class of antiarrhythmic agents. *Chemistry of Natural. Chemistry of Natural Compounds* 1997, 2, 257-270.
3. Karaki H., Ozaki H, Hori M., Nakazawa H., Won R., Sato K. Calcium movements, distribution, and functions in smooth muscle // *Pharmacol. Rev.* - 1997. - V. 49 (2). - P. 157-230.
4. Leung F. P., Yung L. M., Yao X., Laher I., and Huang Y., "Store-operated calcium entry in vascular smooth muscle," *British Journal of Pharmacology*, vol. 153, no. 5, pp. 846-857, 2008.
5. Mirzayeva Yu.T., Sultankhodjaev M.N., Usmanov P.B. Action of diterpenoid alkaloids karakolin and 1-O-benzoylkarakolin on the contractile activity of smooth muscle cells of the rat aorta // *Uzbek Biological Journal.*-2014.-#1.- pp.8-11.
6. Mirzayeva Yu. T., Zaripov A. A., Abrayeva Z. C. The effects of 14-benzoylkarakoline on contractive activity of rat aorta smooth muscle cells // *Евразийский журнал медицинских и естественных наук.* – 2024. – Т. 4. – №. 2 (Special Issue). – С. 383-384.
7. Missiaen L., Parys J. B., De Smedt H., Himpens B., and Casteels R., "Inhibition of inositol trisphosphate-induced calcium release by caffeine is prevented by ATP," *Biochemical Journal*, vol. 300, no. 1, pp. 81-84, 1994.
8. O'Rourke B., Kass D.A., Tomaselli G.F., Käab S., Tunin R., Marbán E. Mechanisms of altered excitation-contraction coupling in canine tachycardia-induced heart failure // *Circ. Res.* - 1999. - V. 84. - P. 562-570.
9. Pogwizd S.M., Qi M., Yuan W., Samarel A.M., Bers D.M. Upregulation of Na<sup>+</sup>/Ca<sup>2+</sup>-exchanger expression and function in an arrhythmogenic rabbit model of heart failure // *Circ. Res.* - 1999. - V. 85. - P. 1009-1019.
10. Sanders K.M. Signal Transduction in Smooth Muscle. Invited Review: Mechanisms of calcium handling in smooth muscles. *J. Appl. Physiol.* 2001., Vol. 91, pp. 1438-1449.
11. Tuliaganov N., Jahangirov F.N., Sadritdinov F.S., Hamdamov I.. "To the pharmacology of some aconitine alkaloids"; 'Pharmacology of plant substances', 1976, pp.76-93, Tashkent.